SYNTHESIS, CHARACTERIZATION, AND INVESTIGATION OF HETEROCYCLIC COMPOUNDS AND THEIR BIOLOGICAL ACTIVITIES

A synopsis for the award of degree Doctorate of Philosophy

Submitted By

SHABEENA PARVEEN

(Roll No: 2165004002)

Enrollment no. – IIMTU/2165190

SUBMITTED TO

Dr. Mojahid Ul-Islam

IIMT COLLEGE OF MEDICAL SCIENCES,

IIMT UNIVERSITY, MEERUT

MAY, 2023

INDEX:

S. No.	Contents	Page No.
1.	INTRODUCTION	1-4
2.	BRIEF REVIEW ON ISOXAZOLE AND BENZIMIDAZOLE	5-6
3.	REVIEW OF LITERATURE	7-10
4.	AIM AND OBJECTIVES	10-11
5.	PLAN OF WORK	12
6.	METHODOLOGY FOR RESEARCH WORK	13-21
6.	REFERENCES	21-24

INTRODUCTION

The synthesis, characterization, and investigation of heterocyclic compounds have garnered significant attention in the discipline of medicinal chemistry becauseof their diverse biological activities. Heterocyclic compounds are organic compounds that contain one or more heteroatoms, such as nitrogen, oxygen, or sulfur, in their ring structure. These compounds exhibit a wide range of pharmacological properties, making them attractive targets for drug discovery and development.

The synthesis of heterocyclic compounds involves the construction of intricate molecular frameworks using various organic reactions and strategies. Through careful design and selection of starting materials, researchers can access a plethora of heterocyclic motifs with diverse chemical and physical properties. The synthetic routes employed often utilize a combination of traditional and modern synthetic methodologies, including multicomponent reactions, transition metal-catalyzed processes, and bio-inspired approaches.

Characterization of heterocyclic compounds is essential to confirm their structural integrity and purity. NMR spectroscopy, X-ray crystallography, infrared (IR) spectroscopy, and are commonly employed for the identification and structural elucidation of synthesized compounds. These techniques provide valuable insights into the connectivity of atoms within the heterocyclic ring, enabling researchers to validate their synthetic strategies.

The biological activities of heterocyclic compounds have attracted considerable attention in the development of new therapeutic agents. By exploring the structure-activity relationship (SAR) of heterocyclic compounds, researchers can optimize their chemical structures to enhance their efficacy, selectivity, and safety profiles.

In this synopsis, we will explore the synthesis, characterization, and biological activities of heterocyclic compounds. We will delve into the various synthetic

strategies employed for the construction of diverse heterocyclic motifs and discuss the analytical techniques used for their characterization. Furthermore, we will investigate the biological activities exhibited by different classes of heterocyclic compounds, highlighting their potential as promising candidates for the development of novel therapeutics.

Overall, the investigation of heterocyclic compounds and their biological activities provides valuable insights into the design and development of new drugs with enhanced pharmacological properties. Understanding the synthesis, characterization, and pharmacological profiles of these compounds contributes to the advancement of medicinal chemistry, offering exciting opportunities for the discovery of novel therapeutics to combat various diseases.

.

BRIEF REVIEW ON ISOXAZOLE AND BENZIMIDAZOLE

Isoxazole and benzimidazole are two classes of heterocyclic compounds that have garnered considerable attention in the field of medicinal chemistry due to their wide array of biological activities and potential therapeutic applications.

Isoxazoles are heterocyclic compounds of five-membered ring containing anti-inflammatory, anticancer, and analgesic activities. The synthesis of isoxazole derivatives can be accomplished through various methods, such as 1,3-dipolar cycloaddition reactions, oxidation of oximes, and cyclization of α,β-unsaturated ketones. The introduction of different substituents on the isoxazole ring can modify their biological activities and enhance their drug-like properties.

.



On the other hand, benzimidazoles are bicyclic heterocycles composed synthesized by condensing o-phenylenediamine with carboxylic acids or their derivatives. Structural modifications in benzimidazoles can be achieved by introducing various substituents on the aromatic ring, thereby influencing their pharmacological profiles and enhancing their selectivity and potency.

The diverse biological activities exhibited by both isoxazoles and benzimidazoles have rendered them attractive targets for drug discovery and development. These compounds often act as inhibitors or modulators of crucial enzymes, receptors, or signaling pathways involved in disease processes. They offer potential therapeutic opportunities in areas such as cancer, infectious diseases, inflammation, neurodegenerative disorders, and metabolic disorders.

Extensive research has been conducted on their synthesis, structural modifications, and characterization to enhance their drug-like properties and optimize their pharmacological profiles. Further research and development in these areas may lead to the discovery of novel drugs with improved efficacy, selectivity, and safety for treating various diseases.



59a 59b 59c 59d 59e

LITERATURE REVIEW

Heterocyclic compounds are a prominent class of organic molecules that have undergone extensive research due to their diverse biological activities and potential therapeutic applications. The synthesis and characterization of these compounds play a vital role in drug discovery and development. This literature review aims to provide an original overview of recent advancements in the synthesis, characterization, and biological activities of heterocyclic compounds.

Synthesis of Heterocyclic Compounds:

Throughout the years, numerous synthetic strategies have been utilized, including classical and modern approaches, to access a wide range of heterocyclic motifs. These methods encompass cyclization reactions, such as

condensation, ring-closing metathesis, and cycloaddition reactions. Additionally, transition metal-catalyzed reactions, like cross-coupling and C-H activation, have emerged as powerful tools for constructing heterocyclic cores. Furthermore, multicomponent reactions have gained significant attention due to their efficiency and ability to explore diverse chemical space.



Characterization Techniques:

Characterizing heterocyclic compounds is crucial to confirm their structural integrity and purity. Various analytical techniques have been employed to elucidate the chemical structure of synthesized compounds. Nuclear magnetic resonance (NMR) spectroscopy, particularly 1H and 13C NMR, provides valuable information about atom connectivity within the heterocyclic ring, enabling the determination of stereochemistry and confirmation of proposed structures. Mass spectrometry (MS) aids in determining the molecular weight and fragmentation pattern of compounds, facilitating their identification. Additionally, techniques such as infrared (IR) spectroscopy, elemental analysis,

and X-ray crystallography contribute to the characterization and confirmation of heterocyclic compounds.

Biological Activities of Heterocyclic Compounds:

For instance, the introduction of different substituents on the heterocyclic ring can significantly influence the biological activity and selectivity of these compounds. Understanding the structure-activity relationship (SAR) is crucial for optimizing their potency, efficacy, and safety profiles. The exploration of heterocyclic scaffolds in drug discovery has led to the development of several successful drugs, such as benzimidazoles, imidazopyridines, and triazoles.



The diverse synthetic methodologies employed have facilitated access to a wide range of heterocyclic motifs, while characterization techniques have enabled the confirmation of their structures. Exploring their biological activities has resulted in the development of novel drugs for various diseases. Further research in this field will continue to uncover the potential of heterocyclic compounds as promising candidates for drug discovery and development, contributing to advancements in medicinal chemistry and healthcare.

AIM AND OBJECTIVES

AIM

To synthesize and Characterization of novel benzimidazole derivatives possessing various biological activities such as Antimicrobial, Anticancer & Antimalarial.

OBJECTIVES

- To investigate and analyze the synthetic methodologies employed for the construction of heterocyclic compounds, including classical and modern approaches, such as cyclization reactions, multicomponent reactions, and transition metal-catalyzed reactions.
- To evaluate the biological activities exhibited by heterocyclic compounds, focusing on their antimicrobial, antiviral, anticancer, anti-inflammatory, and antidiabetic properties.
- To investigate the structure-activity relationship (SAR) of heterocyclic compounds and elucidate the impact of different substituents and modifications on their biological activities.
- To identify potential applications and therapeutic opportunities of heterocyclic compounds in the development of novel drugs for various diseases, including cancer, infectious diseases, inflammation, neurodegenerative disorders, and metabolic disorders.
- To explore the potential of heterocyclic compounds as lead compounds or starting points for further optimization and development of novel drug candidates with improved pharmacological profiles.
- To highlight the importance of interdisciplinary research and collaboration between synthetic chemists, analytical chemists, and biologists in the field of heterocyclic compound synthesis, characterization, and evaluation of their biological activities.

- To contribute to the existing body of knowledge by reviewing and analyzing recent advancements, trends, and challenges in the production, characterization, and biological actions of heterocyclic compounds, providing a comprehensive and up-to-date synopsis in this field.
- To provide a foundation for future research and potential applications of heterocyclic compounds, with a focus on addressing medical needs that are not being met and advancing the field of drug discovery and development..

PLAN OF WORK

Literature survey

↓

Selection of lead molecules

↓

Synthesis of different derivatives of Compound

↓

Method of identification & Characterization

↓

In vitro& In vivo activity

↓

Conclusion

METHODOLOGY FOR RESEARCH WORK

BENZIMIDAZOLE

Scheme- I

Step-I



Where, R-CH_3, Cl



Compound-A

Step-II



Where, R_1 -NO_2,Br, COOH,F

Glacial acetic acid

3-hr



Compound- B

E

Scheme-II



ISOXAZOLE



Selected Examples





Selected Examples



BIOLOGICAL ACTIVITIES

Heterocyclic compounds making them valuable in various fields, including medicine and agriculture. These compounds exhibit diverse pharmacological properties and have shown promise as therapeutic agents. These are some of the significant biological activities associated with heterocyclic compounds:

- Antimicrobial Activity: Heterocyclic compounds have demonstrated potent antimicrobial effects against bacteria, fungi, and other microorganisms. These compounds can inhibit the growth and proliferation of pathogenic microorganisms, offering potential applications in the treatment of infectious diseases.
- Anticancer Activity: Several heterocyclic compounds have exhibited remarkable anticancer activity by interfering with crucial cellular processes, such as cell proliferation, apoptosis, and angiogenesis. These compounds have the potential to be developed as anticancer agents for targeted therapy.
- Antiviral Activity: Heterocyclic compounds have demonstrated considerable potential in the field of antiviral research, exhibiting promising activity against a diverse range of viruses These compounds possess the ability to impede viral replication and diminish viral load, making them valuable candidates for the development of effective antiviral therapies..
- Anti-inflammatory Activity: Heterocyclic compounds possess anti-inflammatory properties and can modulate the immune response. They inhibit the production of pro-inflammatory mediators and enzymes, thus offering potential applications in the treatment of inflammatory diseases.

- Antioxidant Activity: Some heterocyclic compounds exhibit potent antioxidant activity, protecting cells and tissues from oxidative damage. By scavenging free radicals and inhibiting oxidative stress, these compounds may have potential applications in preventing or managing oxidative stress-related disorders.

- Neuroprotection Activity: Specific heterocyclic compounds have shown characteristics of neuroprotection through their influence on neuronal function and preservation. These compounds hold promise as potential therapeutic interventions.

- Antidiabetic Activity: Heterocyclic compounds have been investigated for their potential as antidiabetic agents. They can modulate glucose metabolism, enhance insulin sensitivity, and regulate blood sugar levels, making them promising candidates for the treatment of diabetes.

- Cardiovascular Activity: Some heterocyclic compounds have shown cardiovascular effects by influencing blood pressure, vasodilation, and platelet aggregation. These compounds have potential applications in the management of cardiovascular diseases, such as hypertension and thrombosis.

- Antiparasitic Activity: Heterocyclic compounds exhibit significant antiparasitic activity against various parasites, including protozoa and helminths. They can disrupt parasite growth, reproduction, and survival, making them valuable for the treatment of parasitic infections.

- Analgesic and Anti-inflammatory Activity: Certain heterocyclic compounds possess analgesic properties and can alleviate pain by modulating pain pathways and receptors. Additionally, they exhibit anti-inflammatory effects, reducing inflammation and associated pain.

In summary, heterocyclic compounds exhibit a broad spectrum of biological activities encompassing antimicrobial, anticancer, antiviral, anti-inflammatory, antioxidant, neuroprotective, antidiabetic, cardiovascular, antiparasitic, and analgesic effects. These diverse pharmacological properties underscore the

significance of heterocyclic compounds in the field of drug discovery and development. They offer promising avenues for the creation of novel therapeutic agents targeting a wide range of diseases.

MATERIALS

List of Apparatus & Instruments

Sr. No.	Name(s) of Apparatus and Instruments
1	Beaker
2	Measuring cylinder
3	Pipette
4	Round bottom flask
5	Reflux condenser
6	Glass rod
7	Filter paper
8	China disk
9	Funnel
10	Heating mental
11	Weighing balance
12	Conical flask
13	Test tube
14	Test tube holder
15	Tried stand
16	Magnetic steer
17	Magnetic bead
18	Capillary tube
19	Melting point Apparatus
20	Oven

List of Chemicals

Sr. No.	Name of Chemicals
1	4-chloro-o-phenylenediamine
2	4-methyl-o-phenylenediamine
3	4-fluoro-o-phenylenediamine
4	4-bromo-o-phenylenediamine
5	4-nitro phthalic anhydride4-bromo-o-phenylenediamine
6	4(1-propynyl) phthalicanhydrid
7	Trimettiticanhydrid
8	4-chloro phthalic anhydride
9	4- bromophthalicanhydrid
10	Glacial acetic acid
11	P-toulenesuphoonyl chloride
12	N-ethyl piperzine $C_6H_{14}N_2$
13	4-methoxy benzaldehyde
14	3-chloro benzaldehyde
15	1-methyl piperzine $C_5H_{12}N_2$
16	4-fluoro benzaldehyde
17	2-chloro benzaldehyde
18	1-(2 amino) ethyl piperzine $C_6H_{15}NO_3$
19	2-chloro benzaldehyde
20	4-fluoro benzaldehyde
21	1-(2 amino) ethyl piperzine $C_6H_{15}NO_3$
22	4-nitro benzaldehyde
23	4-chloro benzaldehyde
24	3-bromo benzaldehyde
25	2,O-nitro benzaldehyde

REFERENCES

- Tewari AK and Mishra A: Indian J Chem Sect B: Org ChemIncI Med Chem 2006; 45: 489-493.
- Hoebrecker F: Ber 1872; 5: 920-6.
- Wright JB: January 29, the chemistry of the Benzimidazoles, Research Laboratories, The Upjohn Company, Kalamazoo, Michigan 1951.
- Rathod CP, Rajurkar RM and Thonte SS: Indo American Journal of Pharmaceutical Research 2013; 3(2): 2323-2329.
- Phillip M: J ChemSoc C 1951; 1143-5.
- Hollan G, Samuel L, Ennis B and Hinde R: J ChemSoc C 20-6. Rithe SR, Jagtap RS, Ubarhande SS, 2015. RASA-YAN. J Chem 1967; 8(2): 213-217.
- Saberi A: Iranian Journal of Science & Technology 2015; 39A1: 7-10.
- Suheyla O, Betul F, Canan K and Hakan G: ActaCryst 2002; E58: 1062-1064.
- Mann J, Baron A, Opoku-Boahen Y, Johansson E, Parkinson G, Kelland LR and Neidle S: J Med Chem 2001; 44: 138-144.
- Venkateswarlu Y, Kumar SR and Leelavathi P: Organic and Medicinal Chemistry Letters 2013; 3(7): 2-8.
- Lin S and Yang L: Tetrahedron Letters 2005; 46: 4315-4319.
- Wundt E: The reaction of o-phenylenediamine with formic acid. ChemBer 1878; 11: 826
- Wagner EC: The chemistry of formic acid and its simple derivatives, Org. synthesis 1946; 2: 65.
- Preston PN: Chemistry of heterocyclic compounds benzimidazoles and congeneric tricyclie compounds, Chem Rev 1974; 74: 279.
- Lin XJ: Synthesis crystal structure and photophysical properties of 2-6-bis (benzo[1, 2-d: 5-d] diimidazole -2'-yl-pyridine. Chinese J InorgChem 2004; 20(11): 1316.
- Sridevi C: Designing and biological evaluation of new benzimidazole compounds. ChemSciTra 2013; 2(3): 922.

- Phillips MA: Phillips-Ladenburg benzimidazole synthesis. J ChemSoc 1928; 172.
- Ramanathan V: Synthesis, anti-bacterial, anti-asthmatic and anti-diabetic activities of novel N-substituted 2-(4-styrylphenyl)-1H-benzimidazole and N- substituted-3[4-(1Hbenzimidazole-2-yl)-phenyl]-acrylic acid tert-butyl ester. Arkivoc 2008; 14: 37.
- Sreena K: Synthesis and anthelmintic activity of benzimidazole derivatives. Hygeia 2009; 1(1): 21.
- Roeder H: Benzimidazole and imidazole synthesis. J Org Chem 1941; 6: 25.
- Raut CN: Microwave-assisted sonogashira coupling of novel 2-[6-(arylethynyl) pyridin-3-yl]-1H-benzimidazole derivatives. Arkivoc 2009; 11: 105.
- Niementowski VS: Benzimidazole synthesis. Ber 1897; 30: 3062.
- Rushi T, Surya KD and Richard AG: Journal of Molecular Catalysis A: - Chemical 2006; 245: 8-11.
- Sehyun P, Jaehun J and Eun JCE: J Org Chem 2014; 4148-4154.
- Mobinikhaledi A, Hamta A, Kalhor M and Shariatzadeh M: Iranian Journal of Pharmaceutical Research 2014; 13(1): 95-101.
- Birajdar SS, Hatnapure GD, Keche A P and Kamble VM: Research Journal of Pharmaceutical, Biological and Chemical Sciences 2014; 5(1): 487-493.
- Srinivasulu R, Kumar KR and Satyanarayana PVV: Green and Sustainable Chemistry 2014; 4: 33-37.
- Sehyun P, Jaehun J and Eun JCE: J Org Chem 2014; 4148-4154.
- Vishvanath DP and Ketan PP: International Journal of ChemTech Research 2014; 8(11): 457-465.
- Aniket P, Shantanu DS, Anagha OB and Ajinkya PS: International Journal of ChemTech Research 2015; 8(2): 496-500**.**
- Gupta R. R., Kumar M., Gupta V. Heterocyclic Chemistry, Springer Publication, 1998 1, 13-14.

- Kato T.117. Hosur, M. C.; Talwar, M. B.; Lada, U. V., *Ind. J. Heterocycl. Chem*., 1994, 3, 237-242.
- Ei-Borai, M. A.; Fahmy, M.; Rizk, H., *Ind. J. Heterocycl. Chem*., 1993, 3, 19-24.
- Singh, H.; Yadav, L. D.; Sharma, K. S.; *Ind. J. Chem.*, 1983, 22(B), 1177-1178.
- Boschelli, D. H.; Connor, D. T.; Bornemeier, D. A., *J. Med. Chem.,* 1993, 36, 1802-1810.
- Hui, X. P.; Chang-Hu, C.; Zhang, Z. Y., *Ind. J. Chem.*, 2000, 41(B), 2176-2179.
- Dubey, A. K., Sangwan N. K., *Ind. J. Chem.*, 1994, 33(B), 1034-1047.
- Pattan, S. R.; Rabra, P. A.; Pattan, J. S.; Bukittagar, A. A.; Wakale, V. S.; Musmade, D. S., *Ind. J. Chem.*, 2009, 48(B), 1453-1456.
- Parekh, H.; Patel, P.; Upadhaya, P., *Ind. J. Chem.*, 1996, 35(B), 1062-1066.
- Mulichen, N. D.; Wilson, M. W.; Connor, D. T., *J. Med. Chem*., 1993, 1090-1099.
- Sikder, N.; Bulakh, N. R.; Sikder, A. K., *Ind J. Heterocycl. Chem*., 2002, 12, 29-32.
- Singh, S. P.; Naithani, K.; Batra, H., Ind J. Heterocycl. Chem., 1998, 8, 103-108.
- Sun, X. W.; Liang, H. T.; Zhang, Y., *Ind. J. Chem.*, 1998, 38B, 679-683.
- Hiremath, S. P.; Goudar, N. N., *Ind. J. Chem.*,1982, 21(B), 321-324.
- Datta, M. M.; Goswami, B. N.; Kataky J. C., *J. Ind. Chem. Soc*., 1987, LXIV, 195-198.
- Parekh, O.; Sharma, V.; Batra, H., *Ind. J. Chem*., 1998, 37, 797-800.
- Shaban, M. A. E.; Nasr, A. Z.; El-Badry, S. M., 1991, 4(3), 184-191.
- Gowda, J.; Khadar, A. M. A.; Kalluraya, B.; Kumari, N. S., *Ind. J. Chem*., 2010, 49(B), 1130-1134.
- Ramalingam, R. T.; Deshmukh, A. A.; Sattur, P. B.; Sheth, U. K.; Naik S. R., *J. Ind. Chem. Soc.*, 1981, 58, 269.
- Lazer, E. S.; Farina, P. R.; Oliver, J. T., Possanza, G. J.; Matteo, R., *Agents Action*, 1987, 257.
- Sridhar, S. K.; Ramesh, A., *Bio. Pharm. Bull*., 2001, 24, 1149.
- Scot, F. L.; Lambe, T. M.; Butter, R. N., *J. Chem. Soc. Perkin Trans*, 1972, 1, 269.
- Boschelli, D. H.; Connor, D. T.; Kostlan, C. R.; Kramer, J. B.; Mullican, M. D.; Sircar, J. C., *Eur. Pat.*, 1991, 449211; *Chem. Abstr*., 1992, 116, 6570.

- Omar, F. A., Mahfouz, N. M., Rahman, M. A., *Eur. J. Med. Chem*., 1996, 31, 819-825.
- Palaska, E.; Sahin, G.; Kelicen, P.; Durlu, N. T.; Altinok, G., *Farmaco*, 2002, 57, 101-107.
- Burbuliene, M. M.; Jakubkiene, V.; Mekuskiene, G.; Udrenaite, E.; Smicius, R.; Vainilavicius, P., *Farmaco*, 2004, 59, 767.
- Nargund, I. V. G.; Reddy, G. R. N.; Hariprasad, V., *J. Pharm. Sci*., 2006, 83, 246.